

Dear Patients- In order to make your initial visit more productive, it is necessary that all enclosed papers be <u>read PRIOR to your appointment</u>. PLEASE if you have questions; call the office **before** your initial appointment.

** PLEASE DO NOT WEAR... <u>PERFUMES</u> or <u>SCENTED COLOGNES</u>. It may irritate or harm our patients with allergies or asthma. If patient has long hair, please wear or bring something to tie it back with. Thank you.

1. DO NOT TAKE ANTIHISTAMINES PRIOR TO VISIT.

Check cold preparations for antihistamines. These are some of the major brands to look for:

*MUST BE OFF 7 DAYS PRIOR TO VISIT:

ALL VITAMINS, ALL SUPPLEMENTS

VITAMIN C, significant amount of Orange Juice

Allegra/ Allegra-D (Fexofenadine) Xyzal (Levocetirizine)

Zyrtec/ Zyrtec-D (Cetirizine) Claritin/Claritin-D/Alavert (Loratadine)

Clarinex/ Clarinex-D (Desloratadine) Atarax/ Vistaril (Hydroxizine)

Semprex-D Tagament (Cimetidine)
Pepcid (Famotidine) Axid (Nizatidine)
Zantac (Ranitidine) Probiotics

Zantac (Ranitidine) Cyproheptadine (Periactin)

MUST BE OFF 3 DAYS PRIOR TO VISIT:

Dimetapp Cyclizine
Tavist/ Tavist D Doxylamine
Patanase, Patanol, Pataday (Olopatadine) Promethazine

Astelin, Astepro, Dymista (Azelastine)

Benadryl/Diphenhydramine

Chlorpheniramine

Dramamine (Dimenhydrinate

- **2**. Possible **BEE / VENOM** reaction patients, if this is the **only** problem you are coming for, you **do not have to stop antihistamines.** You will be tested another day.
- **3**. **ANTIBIOTIC REACTION-** If you are coming for **only** an antibiotic reaction, you do not have to stop antihistamines. You will be tested another day.
- 4. If you are scheduled for a breathing test (Pulmonary Function Test)
- *DO NOT TAKE any oral, inhaled or nebulized bronchodilators the day of your appointment.

Common types are... Albuterol (Proventil, Ventolin or ProAir), Xopenex (Levalbuterol), Asmanex

Spiriva, Serevent, Breztri, Breo, Trelegy and Airduo, Symbicort, Xopenex, Advair/(Fluticasone Salmeterol), Atrovent (Ipratropium bromide), Combivent & Combivent Respimat, Airsupra Fluticasone

- *If you are unsure, please feel free to call.
- *However, use your inhaler or nebulizer if you are experiencing shortness of breath, difficulty breathing, wheezing, or a severe cough.
- 5. CONTINUE TAKING any Theophylline's, Oral or Inhaled Steroids, Singulair, Accolate.
- **6. BRING** complete list of **all medications**; if possible bring recent Blood work, Sinus or Chest X-Rays.

YOUR INITIAL VISIT ranges in time from 2–4 hours and includes Consultation, Physical Exam, Allergy Testing and for some patients, Pulmonary Function Testing. Additional testing may be required and completed another day. Should you fail to follow the above directions and cancel the NEW patient appointment without a 24-hour notice, a \$150 fee will be assessed. In addition, if you do not stop medications as instructed and present for appointment, you will have a Consult only and will be directed to return for another appointment.

SIGNATURE OF PATIENT/GUARDIAN	PATIENT NAME	DATE